

Production of the π^- , π^0 in the reaction $d(e, pp)e'\pi^-\pi^0$ ¹.

A.Yu. Loginov^a, D.M. Nikolenko^b, A.V. Osipov^a, I.A. Rachek^b,
 A.A. Sidorov^a,

V.N. Stibunov^a, D.K. Toporkov^b

^a*Nuclear Physics Institute at Tomsk Polytechnic University, Tomsk, Russia*

^b*Budker Institute of Nuclear Physics, Novosibirsk, Russia*

Talk presented at the 15th International Conference
 on Particles and Nuclei, 10-16 June, 1999,
 Uppsala University, Uppsala, Sweden.

Abstract

Recently, the differential yield and target asymmetry of the reaction $d(e, pp)e'\pi^-$ on the tensor polarized deuterium target have been presented in [1, 2], where the experimental procedure and basic assumption were described. A precise treatment of this reaction needs to investigate the reaction channels with an additional neutral pion production as a background to the leading reaction of a single π^- production.

We considered for this background process the pole mechanism and the two body mechanism. In the latter the Δ -isobar and the pion are produced on one nucleon followed by the absorption of the pion on the second nucleon producing another Δ -isobar. The final proton-pion systems come from the decay of the Δ^+ and Δ^0 .

In this work we estimated the contribution of the reaction with additional π^0 production into whole yield for the different polarization states of the deuteron.

In a study of the $d(e, pp)X$ - reaction at electron energy $2GeV$, the momenta of both protons from 0.3 to 0.7 GeV/c and proton emission angles $64^\circ - 83^\circ$ approximately 900 proton proton coincidences were analyzed [1, 2]. The cross section of the process initiated by electrons was expressed in terms of the cross section of a reaction induced by virtual photons:

$$\gamma^* d \rightarrow ppX. \quad (1)$$

In the forward scattering approximation a virtual photon can be replaced by a real photon. Assuming that only a single π^- were produced, the photon energy

¹ This work was supported by Russian Foundation for Basic Research N 98-02-17993, N 98-02-17949 and by grant N 96-0424 from INTAS.

can be calculated from energy and momentum balance using momenta of the both protons, defined in the experiment. One can say that experimental distributions essentially exceed the theoretical estimations made in the one particle frameworks. The background event contribution that it does not take into account could be a source of this difference. Since there is no additional equation, one cannot distinguish the background events with one and more neutral particles. Although an admixture of the background process with additional π^0 according to the experimental estimations is small, we decided to calculate it directly. It should be noted that the maximum value of the missing mass in our experiment[1]:

$$M_{mis} = \sqrt{(M_d + E_e - E_1 - E_2)^2 - (\vec{p}_e - \vec{p}_1 - \vec{p}_2)^2} \quad (2)$$

is obviously less than the ρ - meson mass, so background from ρ - production is absent.

We have studied the background process $\gamma d \rightarrow pp\pi^-\pi^0$ as two different possibilities. The first is the production of the $\pi^-\pi^0$ on the neutron – let us call it the pole contribution and the two particle process of the $\Delta^+ \Delta^0$ production when the final πp systems comes from the decay of Δ according to the diagrams depicted in Fig.1. These channels do not exhaust all possibilities. The model of the production of two pions on the nucleon and the deuteron was developed in the works [3, 4]. We used it for the tensor polarized deuteron and large momenta of the decayed protons in our isotopic channel.

The matrix elements of the pole and double Δ reaction mechanisms are:

$$\begin{aligned} iM_{M_d \lambda}^{(pole)m_1 m_2} = & \frac{\sqrt{2}}{3} e \left(\frac{f^*}{\mu} \right)^2 \sum_{m'_1} \chi_{m_1}^\dagger \left(\frac{2}{3} \vec{q}_- \cdot \vec{\epsilon}_\lambda - \frac{i}{3} \vec{q}_- \cdot [\vec{\epsilon}_\lambda \vec{\sigma}] \right) \chi_{m'_1} \times \\ & \times G_\Delta(p_1 + q_0) \Psi_{m'_1 m_2, M_d}(\vec{p}_2) - \\ & - \frac{\sqrt{2}}{3} e \left(\frac{f^*}{\mu} \right)^2 \sum_{m'_1} \chi_{m_2}^\dagger \left(\frac{2}{3} \vec{q}_- \cdot \vec{\epsilon}_\lambda - \frac{i}{3} \vec{q}_- \cdot [\vec{\epsilon}_\lambda \vec{\sigma}] \right) \chi_{m'_1} \times \\ & \times G_\Delta(p_2 + q_0) \Psi_{m'_1 m_1, M_d}(\vec{p}_1) \end{aligned} \quad (3)$$

$$\begin{aligned} iM_{M_d \lambda}^{(delta)m_1 m_2} = & i \frac{\sqrt{2}}{9} e \left(\frac{f^*}{\mu} \right)^4 \int \frac{d^3 n'}{(2\pi)^3} \sum_{m'_1 m'_2} \chi_{m_1}^\dagger \left(\frac{2}{3} \vec{q}_- \cdot \vec{\epsilon}_\lambda - \frac{i}{3} \vec{q}_- \cdot [\vec{\epsilon}_\lambda \vec{\sigma}] \right) \chi_{m'_1} \times \\ & \times \chi_{m'_2}^\dagger \left(\frac{2}{3} \vec{q}_0 \cdot \vec{q}' - \frac{i}{3} \vec{q}_0 \cdot [\vec{q}' \vec{\sigma}] \right) \chi_{m'_2} F^2(q'^2) G_\Delta(p_1 + q_-) \times \end{aligned}$$




Figure 1:

$$\begin{aligned}
& \times G_{\Delta}(p_2 + q_0) \frac{1}{q'^2 - \mu^2 + i\epsilon} \Psi_{m'_1 m'_2, M_d}(\vec{n}') + \\
& + i \frac{\sqrt{2}}{9} e \left(\frac{f^*}{\mu} \right)^4 \int \frac{d^3 p'}{(2\pi)^3} \sum_{m'_1 m'_2} \chi_{m'_1}^\dagger \left(\frac{2}{3} \vec{q}_- \cdot \vec{q}' - \frac{i}{3} [\vec{q}_- \cdot \vec{\sigma}] \right) \chi_{m'_1} \times \\
& \times \chi_{m'_2}^\dagger \left(\frac{2}{3} \vec{q}_0 \cdot \vec{\epsilon}_\lambda - \frac{i}{3} \vec{q}_0 \cdot [\vec{\epsilon}_\lambda \cdot \vec{\sigma}] \right) \chi_{m'_2} F^2(q'^2) G_{\Delta}(p_1 + q_-) \times \\
& \times G_{\Delta}(p_2 + q_0) \frac{1}{q'^2 - \mu^2 + i\epsilon} \Psi_{m'_1 m'_2, M_d}(\vec{p}') - \\
& - (1 \leftrightarrow 2) .
\end{aligned} \tag{4}$$

where M_d , λ , m_1 , m_2 are spin projection of the initial and final particles; e is the electron charge; μ , M_{Δ} are the meson and Δ masses; p_1 , p_2 , n' , p' , q_0 , q_- are four momenta of the nucleons and pions according to diagrams depicted in Fig.1. Coupling constant $f^*/4\pi = 0.36$. ϵ_λ is the photon polarization vector (in Coulomb gauge $\epsilon_0 = 0$, $\vec{\epsilon} \cdot \vec{k} = 0$), $F(q^2)$ a monopole form factor with $\Lambda = 1.3 \text{ GeV}$,

$\Psi_{m_1 m_2, M_d}(\vec{p})$ is the deuteron relative wave function in momentum space. For the Δ propagator we take the formula:

$$G_{\Delta}(p) = \frac{1}{\sqrt{s} - m_{\Delta} + \frac{i}{2}\Gamma_{\Delta}(s)} \frac{M_{\Delta}}{E_{\Delta}(\vec{p})} . \quad (5)$$

The two proton yield from the reaction $ed \rightarrow pp\pi^{-}\pi^0 e'$:

$$Y = L \int f(\omega) \frac{d^8 \sigma_{\gamma^*}(\omega)}{d^3 p_1 d^3 p_2 d\Omega_q} d^3 p_1 d^3 p_2 d\Omega_q , \quad (6)$$

where L -experimental luminosity, Ω_q is solid angle of the relative momentum of the negative and neutral pions. We have selected (θ_i, ϕ_i, p_i) of the protons from our experimental regions [1] and integrated the differential cross section of the background process using the Monte Carlo method.




Figure 2: The total cross section of the $\pi^{-}\pi^0$ photoproduction on the deuteron target as a function of the photon energy (solid line - pole mechanism, dashed line - $\Delta^+\Delta^0$ production).




Figure 3: The yield of the $\pi^-\pi^0$ production the pole mechanism as a function of the angle between proton detector axis and photon beam direction.




Figure 4: The yield of the $\pi^-\pi^0$ production for double Δ mechanism as a function of the angle between proton detector axis and photon beam direction.

Both pions can fly in 4π geometry, so we have integrated differential cross section by $d\Omega_q$ over the full solid angle. The pion energies was determined by the conservation laws. The energy spectrum of the virtual photons has the form presented in [5].

For the checking we estimated the total cross section of the $\pi^-\pi^0$ photoproduction on the deuteron target by the same method for both mechanisms and presented them in Fig.2. The total cross section as a function of the photon energy for the studied isotopic channel agrees with the results of [4] for another isotopic channel if the Δ^+, Δ^0 decay would be considered.

The results of the estimation of Y for the two reaction mechanisms are presented in Fig.3 and Fig.4 for two signes of the deuteron target tensor polarization: ($P_{zz} = 0.573$ – solid and dashed lines, $P_{zz} = -0.573$ – dotted and dash-dotted lines) and for two directions of the magnetic field: (solid and dotted lines – $\vec{H}$ in the plane of the proton detector axes, dashed and dash-dotted lines – $\vec{H}$ perpendicular to that plane).

These four curves is shown only in illustrative purposes as at our experimental

conditions they cannot be distinguished. These background events are presented as a function of the angle between the proton detector axis and photon beam direction when the proton detectors are placed symmetrically about this direction. One can see that the total cross section, calculated with pole matrix elements, drastically exceeds that for the second mechanism. In order to estimate the contributions of both mechanisms into the calculated yield at the experimental region [1], it is necessary to put the angle θ in Figs.3, 4 equal to 75° . So the background yield amounts to about three events. It is too small to be represented together with experimental and theoretical distributions of the proton-proton events from the main process [1]. Indeed, the whole experimental yield is about 440 events at one system of the proton-proton registration and the theoretical yield, calculated in a spectator model, is about 250 events.

A assumption about the appreciable contribution of these background events turned out to be unfounded. A number of theoretical models were used for explanation of experimental data [1]. The approaches based on the traditional concepts seem to be unable to reproduce all features of the proton-proton production from the deuteron [1, 2].

References

- [1] *A.Yu. Loginov, A.V. Osipov, A.A. Sidorov, V.N. Stibunov et al.* // Pis'ma Zh. Eksp. Teor. Fiz. 1998. V.67 N 10 P.730; JETP Letters 1998. V.67 N 10 P.770
- [2] *V.N. Stibunov et al.* // Few-Body Systems Suppl. 10. 1999. P.507
- [3] *J.A. Gómez Tejedor, E. Oset* // Nucl. Phys. 1994. V.A571 P.667
- [4] *J.A. Gómez Tejedor, E. Oset, H. Toki* // Phys. Lett. 1995. V.B346 P.240
- [5] *R.H. Dalitz, D.R. Yennie* // Phys. Rev. 1957. V. 105 P. 1598